

UC Davis

UC Davis Previously Published Works

Title

Neuron numbers increase in the human amygdala from birth to adulthood, but not in autism.

Permalink

<https://escholarship.org/uc/item/1gm452hh>

Journal

Proceedings of the National Academy of Sciences of the United States of America, 115(14)

ISSN

0027-8424

Authors

Avino, Thomas A
Barger, Nicole
Vargas, Martha V
et al.

Publication Date

2018-04-01

DOI

10.1073/pnas.1801912115

Peer reviewed

Neuron numbers increase in the human amygdala from birth to adulthood, but not in autism

Thomas A. Avino^a, Nicole Barger^a, Martha V. Vargas^a, Erin L. Carlson^a, David G. Amaral^{a,b,c}, Melissa D. Bauman^{a,b}, and Cynthia M. Schumann^{a,1}

^aDepartment of Psychiatry and Behavioral Sciences, UC Davis MIND Institute, School of Medicine, University of California, Davis, Sacramento, CA 95817;

^bCalifornia National Primate Research Center, University of California, Davis, CA 95616; and ^cCenter for Neuroscience, University of California, Davis, CA 95618

Edited by Joseph E. LeDoux, New York University, New York, NY, and approved March 1, 2018 (received for review February 12, 2018)

Remarkably little is known about the postnatal cellular development of the human amygdala. It plays a central role in mediating emotional behavior and has an unusually protracted development well into adulthood, increasing in size by 40% from youth to adulthood. Variation from this typical neurodevelopmental trajectory could have profound implications on normal emotional development. We report the results of a stereological analysis of the number of neurons in amygdala nuclei of 52 human brains ranging from 2 to 48 years of age [24 neurotypical and 28 autism spectrum disorder (ASD)]. In neurotypical development, the number of mature neurons in the basal and accessory basal nuclei increases from childhood to adulthood, coinciding with a decrease of immature neurons within the paralamina nucleus. Individuals with ASD, in contrast, show an initial excess of amygdala neurons during childhood, followed by a reduction in adulthood across nuclei. We propose that there is a long-term contribution of mature neurons from the paralamina nucleus to other nuclei of the neurotypical human amygdala and that this growth trajectory may be altered in ASD, potentially underlying the volumetric changes detected in ASD and other neurodevelopmental or neuropsychiatric disorders.

autism | amygdala | stereology | neuroanatomy | neuronal maturation

The human amygdala comprises a cluster of 13 nuclei in the rostral temporal lobe which play a critical role in fear, emotion, and social behavior (1–3). The typical human and nonhuman primate amygdala undergoes a remarkable 40% volumetric growth into early adulthood, despite little growth of the cerebral cortex (4, 5). Prolonged amygdala maturation likely underlies the increasing functional integration of this structure as it modulates responses to an ever-changing environment. Early perturbations in amygdala cellular development could lead to a cascade of maladaptive neurodevelopmental events that affect the entire trajectory of maturation. Revealing the underlying neurobiology of this prolonged growth is critical not only for a fundamental understanding of neurotypical development but also for pinpointing when these processes deviate in disorders such as autism spectrum disorder (ASD) and other neuropsychiatric disorders (6, 7).

Although the primate amygdala is formed early in gestation and is well developed at birth (8–11), structural and functional changes extend well into adulthood (12, 13). A number of factors likely contribute to the dramatic increase in postnatal amygdala volume, including dendritic enlargement, synaptogenesis, and gliogenesis. We propose that other neuronal factors may contribute, such as (i) the maturation of a large population of immature neurons within the paralamina nucleus (6, 14–17) and/or (ii) the migration of postnatally generated neurons (18, 19). Immature neurons have been identified in both monkey and human amygdala using immunohistochemistry with protein markers such as doublecortin (DCX) and B cell lymphoma 2 (bcl-2) (14, 16–18, 20, 21). The protracted maturational trajectory of the amygdala well beyond the perinatal period allows it to continually be shaped by external stimuli. In fact, the immature neurons within the paralamina nucleus may develop in an activity-dependent manner (20, 22). This lengthened maturational

process, however, may also make the amygdala more susceptible to developmental or environmental insults.

ASD is characterized by impairments in social communication combined with restricted interests and behaviors. Alterations in amygdala growth can be detected as early as 2 y of age (23–26) and persist into late childhood (5, 27). The severity of the individual's social and communicative symptoms positively correlates with amygdala enlargement, suggesting a potential structure–function relationship (23). Individuals with ASD also show atypical amygdala activation during socioemotional tasks (28, 29). Microanatomical alterations to the cellular structure of the amygdala were first noted by Bauman and Kemper (30) and subsequently by Schumann and Amaral (31) and Wegiel et al. (32). These studies found a general reduction of neurons in the amygdala of adults with ASD. However, an examination of younger subjects with either neurotypical development or ASD has not yet been performed. The present study aimed to carry out a large systematic evaluation of the developmental trajectory of neuron number from youth to adulthood in the human amygdala in both neurotypical individuals and in those diagnosed with ASD. In addition, we examined the presence of immature neurons in the amygdala and evaluated whether differences in this population across the life span may contribute to the gradual decreases in neuron number we have observed in our previous studies of adults with ASD.

Results

Case Information. Subject-specific information from all 52 subjects is presented in [Table S1](#). Briefly, the neurotypical-subject (i.e., NT) group contained 24 subjects (mean age, 20.17 ± 13.28 y; range, 2 to 48 y), 5 of whom were female. The ASD group

Significance

We demonstrate that the number of mature neurons in the human amygdala increases from childhood into adulthood. This trajectory may be due to the incorporation of immature neurons from the paralamina nucleus in the ventral amygdala. In contrast, individuals with autism spectrum disorder (ASD) show an initial excess of mature neurons followed by a decline into adulthood. Our results suggest a degenerative component in ASD and highlight the need for a more comprehensive understanding of the protracted cellular development of the human amygdala for multiple psychiatric disorders.

Author contributions: T.A.A., N.B., D.G.A., and C.M.S. designed research; T.A.A., N.B., M.V.V., E.L.C., and C.M.S. performed research; T.A.A., N.B., M.D.B., and C.M.S. analyzed data; and T.A.A., N.B., D.G.A., M.D.B., and C.M.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](#).

¹To whom correspondence should be addressed. Email: cschumann@ucdavis.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1801912115/-DCSupplemental.

Published online March 20, 2018.




Fig. 1. Mature neuron number across amygdala nuclei between ASD and NT subjects. Young subjects with ASD show an increased number of mature neurons relative to NT subjects (in total amygdala, basal, accessory basal, and central nuclei). By adulthood, the number of mature neurons in ASD is well below the adult NT average in every nucleus examined (17%). Error bars ± 1 SEM. When considering the monkey basal + paralaminar nuclei (6) as a single unit (as we have done with the human basal nucleus, *Lower Right*), there is a comparable increase of mature neurons ($\sim 32\%$) across life between the two species.


neurons in the amygdala of ASD cases appears to decline from childhood to adulthood by about 17% and is significantly reduced in all regions we examined. Third, in the amygdala from both neurotypical and ASD cases, there is a large pool of bcl-2+ immature neurons in the paralaminar nucleus, which declines with age from childhood into adulthood.

Together, our data indicate that as the number of mature neurons increases in the neurotypical amygdala across life, the number of immature neurons in the paralaminar nucleus decreases. Therefore, it is possible that a gradual maturation and migration of immature neurons from the paralaminar nucleus contributes to the increasing number of mature neurons in the neurotypical amygdala, particularly in the basal and accessory basal nuclei. Individuals with ASD demonstrate a similar decrease in immature neurons in the paralaminar nucleus with age, yet the net number of mature neurons does not increase with age. Rather, cases of ASD show a decline in mature neuron number across the life span, suggesting a substantial loss of amygdala neurons throughout life in individuals with ASD. We discuss the rationale for this hypothesis below.

Typical Pattern of Amygdala Neuronal Development from Youth Through Adulthood. Structural MRI studies have demonstrated that the amygdala in neurotypical individuals continues to grow in size by $\sim 40\%$ throughout adolescence into adulthood. This is in contrast to most of the cerebral cortex, which ceases to grow appreciably after 6 y of age (4, 5, 12, 13). There was little or no understanding of the neurobiological substrate for this postnatal

increase in volumetric growth of the human amygdala. We now report that the number of mature neurons in the amygdala increases by $\sim 11\%$ from early childhood into adulthood. However, the extent of neuronal increase varies across the different subregions of the amygdala, where it was greatest in the basal nucleus ($\sim 30\%$), more modest in the accessory basal nucleus ($\sim 17\%$), and much less so in the lateral nucleus ($\sim 3\%$).

One hypothesis to account for this increase in neuron number is the slow maturation and migration of a large pool of immature neurons within in the paralaminar nucleus produced prenatally. Here, we document a marked reduction in the number of immature (bcl-2+) cells in the human paralaminar nucleus, while mature basal nucleus neuron numbers increase from youth to adulthood. Bcl-2 immunoreactivity has been used to identify immature neurons in both human (16, 17) and nonhuman primate amygdala (6, 15, 18). A variety of other immature and migratory neuronal markers, including DCX, polysialylated-neural cell adhesion molecule, and class III β -tubulin (14), confirm their presence. Although the majority of bcl-2+ and DCX+ cells are found in the paralaminar nucleus ventral to the parvocellular portion of the basal nucleus, the paralaminar nucleus has a broad expanse, wrapping around the rostral amygdala. We also observed bcl-2+ and DCX+ cells in the ventral lateral nucleus, periamygdaloid cortex, and intercalated islands (15–17, 21, 33). Our data and others' indicate that immature bcl-2 immunoreactive cells likely undergo a protracted maturational period well into adulthood (14), with numbers declining during



PNAS | April 3, 2018 | vol. 115 | no. 14 | 3713

Another postnatal factor that may explain the increase of mature neuron number in the amygdala of individuals with ASD involves the activity-dependent maturation of immature amygdala neurons. Previous work has shown that neurons within the paralamina nucleus may mature in an experience-dependent way (22). A recent report demonstrated that hippocampal-lesioned animals show an increased number of mature and immature neurons in the amygdala, suggesting that the lesions might not only drive neuronal differentiation but also stimulate the production of immature neurons in the nearby subventricular zone (20). It is possible that the hyperactivity of the amygdala frequently observed in individuals with ASD is in some way comparable to the stimulatory effects of these hippocampal lesions. In both instances, insults to the development and connectivity of the amygdala could conceivably lead to excessive neuronal differentiation and production.

Adolescence to adulthood—ASD. During the time when the number of mature neurons within the amygdala is gradually increasing in neurotypical development, adolescents with ASD begin showing a reduction of neurons. The substantial decline of neuron numbers by adulthood in ASD may be driven by prolonged hyperactivation of the amygdala throughout life. There is some support for this hypothesis in other disorders, such as depression, where prolonged overactivation could result in decreases to amygdala and hippocampal volume, potentially through excessive glucocorticoid activity leading to excitotoxic neuron loss (39, 40). Similarly, this excitotoxicity may be occurring in ASD, which frequently cooccurs with anxiety such that by adulthood, the number of amygdala neurons in ASD is reduced relative to both adult-aged neurotypical and pediatric-aged ASD individuals. This excitatory/inhibitory imbalance of the amygdala in ASD may be caused by a number of factors, including a lack of habituation to sensory stimuli (41, 42), a lack of inhibitory control from frontal cortical areas, and changes to cellular and synaptic function. It is also plausible that the decline of neuron numbers in the amygdala of individuals with ASD may also be influenced by glia–neuron interactions or other immune factors. Specifically, microglia have been shown to regulate cell proliferation during the late stages of neurogenesis through phagocytosis of neural progenitors (43). Their role continues in the mature brain where they are responsible for the removal of apoptotic cells. Abnormal microglial presence and activation has been reported in ASD in the dorsolateral prefrontal cortex (44) and in the amygdala (45); however, these alterations were present only in a subset of ASD cases and unrelated to decreases in neuron number.

Together, our results suggest that there is an atypical developmental trajectory in the amygdala of individuals with ASD that results in substantial neuron loss by adulthood. Our findings are in accordance with other neuroanatomical studies that have observed increased numbers of prefrontal cortical neurons in children with ASD (36), as well as reduced neuron numbers in the fusiform gyrus and lateral nucleus of the amygdala (32, 46). However, the findings presented here may not be specific to autism and have extensive implications for a number of neurodevelopmental and neuropsychiatric disorders (7). Amygdala dysfunction has traditionally been implicated in a broad range of neuropsychiatric illnesses, including anxiety, mood disorders, posttraumatic stress disorder, and schizophrenia (47, 48). However, the precise contribution of amygdala dysfunction in these diseases and the developmental time course are not well understood. Future studies will need to address the origin, timing, plasticity, and neuronal fate of these immature neurons to further elucidate their role in neurological diseases and how they may be used as a potential therapeutic target for mental health.

Conclusion

We found that the number of mature neurons in the typical human amygdala increases over a protracted postnatal period, extending into at least late adolescence. We propose that this is due to the maturation and migration of immature neurons that are located initially in the paralamina nucleus of the amygdala. The stimulus for this slow accretion of mature neurons is

currently unclear but appears to be a unique feature of the postnatal development of the amygdala. The trajectory of neuronal development in the amygdala of individuals who had ASD during life drastically deviates from the typical developmental trajectory. There are ~11% more neurons in the amygdala in very young individuals with ASD, but ~20% fewer neurons in adults with ASD. This altered growth pattern may lead to altered amygdala function, manifesting in increased anxiety and further contributing to social impairments. Hyperactivity of the amygdala may also lead to neuron loss through excitotoxicity, especially in instances where the amygdala and its neurons are particularly vulnerable to stress. Understanding the regulation of neuron number at a fundamental level in this brain region will be important for interpreting the normal role of the human amygdala during life and may provide insight into some of the neural disturbances that contribute to the behavioral pathology of multiple neurodevelopmental and psychiatric disorders.

Experimental Procedures

Brain Samples. Tissue series from 52 individuals, aged 2 to 48 y at death, were used for stereological quantification and contained the entire rostrocaudal extent of the amygdala (28 ASD, 24 NT; Table S1). All subjects were included from two cohorts, which differed in tissue-processing protocol. Cohort 1 included 19 ASD and 15 neurotypical brains processed in our laboratory obtained from the NIH NeuroBioBank or Autism BrainNet (formerly Autism Tissue Program). Partial data and diagnostic information from a subset of cohort 1 ($n = 19$) have previously been published by Schumann and Amaral (31). Cohort 2 included nine ASD and nine neurotypical brains from the Autism Celloidin Library distributed by Autism BrainNet. Cohorts 1 and 2 were combined for neuron number estimates and separated for cell volumetric analyses to avoid potential confounding factors due to variation in processing protocols. This study was exempt from Institutional Review Board approval.

Tissue Processing. Detailed descriptions of tissue processing are available in previous studies from cohort 1 (31) and cohort 2 (49, 50) and in [SI Experimental Procedures](#).

Stereological Design. The amygdala and its lateral, basal, accessory basal, and central nuclei (Fig. S2) were delineated using previously published criteria (51). The paralamina nucleus was identified using adjacent Nissl-stained sections. We sampled mature (Nissl) and immature (bcl-2+) neurons (100 \times objective, N.A. 1.3) through the entire structure of interest to derive




Fig. 3. A methodological summary of our stereological approach. The entire rostrocaudal extent of the amygdala is sectioned. For experiment 1, we used Nissl-stained sections at a 1/5 sampling interval, and for experiment 2, we used alternating bcl-2-stained sections at a 1/10 sampling interval from the same brains. Each section has a virtual grid overlaid on top which designates physical locations to place sampling boxes. The numbers of objects are counted in each sampling box, and an estimate is extrapolated based on the size of the sampling box, the density of the sampling grid, the number of sections examined, and tissue thickness. Stereological parameters are presented in Table S2. (Scale bars: Left four, 2 mm; Right two, 10 μm.)

numerical estimates using the optical fractionator as in our prior publications (52, 53). Stereological design is summarized in Fig. 3. Sampling parameters are presented in Table S2.

Statistical Analysis. Subjects were classified a priori into three age groups: pediatric [2 to 13 y old; $n = 17$ (10 ASD, 7 NT)]; adolescent [14 to 20 y old; $n = 11$ (6 ASD, 5 NT)]; and adult [21+ y old; $n = 19$ (9 ASD, 10 NT)]. Univariate two-way ANOVA was used to test for effects of age group and diagnosis on neuron number, region volume, and neuron somal or nuclear volume within each amygdala nuclei. Significant age group \times diagnosis interactions were further analyzed using a Fisher LSD post hoc test. The data met parametric assumptions for normality using a Shapiro–Wilk test ($P > 0.05$), and for homogeneity of variance using Levene's test for equality of variance ($P > 0.05$).

- Adolphs R (2010) What does the amygdala contribute to social cognition? *Ann N Y Acad Sci* 1191:42–61.
- Fox AS, Oler JA, Tromp PM, Fudge JL, Kalin NH (2015) Extending the amygdala in theories of threat processing. *Trends Neurosci* 38:319–329.
- Amaral DG (2002) The primate amygdala and the neurobiology of social behavior: Implications for understanding social anxiety. *Biol Psychiatry* 51:11–17.
- Giedd JN, et al. (1996) Quantitative MRI of the temporal lobe, amygdala, and hippocampus in normal human development: Ages 4–18 years. *J Comp Neurol* 366:223–230.
- Schumann CM, et al. (2004) The amygdala is enlarged in children but not adolescents with autism; the hippocampus is enlarged at all ages. *J Neurosci* 24:6392–6401.
- Chareyron LJ, Lavenex PB, Amaral DG, Lavenex P (2012) Postnatal development of the amygdala: A stereological study in macaque monkeys. *J Comp Neurol* 520:1965–1984.
- Schumann CM, Bauman MD, Amaral DG (2011) Abnormal structure or function of the amygdala is a common component of neurodevelopmental disorders. *Neuropsychologia* 49:745–759.
- Humphrey T (1968) The development of the human amygdala during early embryonic life. *J Comp Neurol* 132:135–165.
- Humphrey T (1972) The development of the human amygdaloid complex. *The Neurobiology of the Amygdala: The Proceedings of a Symposium on the Neurobiology of the Amygdala*, Bar Harbor, Maine, June 6–17, 1971 (Springer, Boston), pp 21–80.
- Müller F, O'Rahilly R (2006) The amygdaloid complex and the medial and lateral ventricular eminences in staged human embryos. *J Anat* 208:547–564.
- Ulfing N, Setzer M, Bohl J (2003) Ontogeny of the human amygdala. *Ann N Y Acad Sci* 985:22–33.
- Østby Y, et al. (2009) Heterogeneity in subcortical brain development: A structural magnetic resonance imaging study of brain maturation from 8 to 30 years. *J Neurosci* 29:11772–11782.
- Uematsu A, et al. (2012) Developmental trajectories of amygdala and hippocampus from infancy to early adulthood in healthy individuals. *PLoS One* 7:e46970.
- deCampo DM, Fudge JL (2012) Where and what is the paralaminar nucleus? A review on a unique and frequently overlooked area of the primate amygdala. *Neurosci Biobehav Rev* 36:520–535.
- Fudge JL (2004) Bcl-2 immunoreactive neurons are differentially distributed in subregions of the amygdala and hippocampus of the adult macaque. *Neuroscience* 127:539–556.
- Marti-Mengual U, Varela E, Crespo C, Blasco-Ibáñez JM, Nacher J (2013) Cells expressing markers of immature neurons in the amygdala of adult humans. *Eur J Neurosci* 37:10–22.
- Yachnis AT, Roper SN, Love A, Fancey JT, Muir D (2000) Bcl-2 immunoreactive cells with immature neuronal phenotype exist in the nonepileptic adult human brain. *J Neuropathol Exp Neurol* 59:113–119.
- Bernier PJ, Bedard A, Vinet J, Levesque M, Parent A (2002) Newly generated neurons in the amygdala and adjoining cortex of adult primates. *Proc Natl Acad Sci USA* 99:11464–11469.
- Gould E, Reeves AJ, Graziano MS, Gross CG (1999) Neurogenesis in the neocortex of adult primates. *Science* 286:548–552.
- Chareyron LJ, Amaral DG, Lavenex P (2016) Selective lesion of the hippocampus increases the differentiation of immature neurons in the monkey amygdala. *Proc Natl Acad Sci USA* 113:14420–14425.
- Bernier PJ, Parent A (1998) Bcl-2 protein as a marker of neuronal immaturity in postnatal primate brain. *J Neurosci* 18:2486–2497.
- Fudge JL, deCampo DM, Becoats KT (2012) Revisiting the hippocampal-amygdala pathway in primates: Association with immature-appearing neurons. *Neuroscience* 212:104–119.
- Schumann CM, Barnes CC, Lord C, Courchesne E (2009) Amygdala enlargement in toddlers with autism related to severity of social and communication impairments. *Biol Psychiatry* 66:942–949.
- Kim JE, et al. (2010) Laterobasal amygdalar enlargement in 6- to 7-year-old children with autism spectrum disorder. *Arch Gen Psychiatry* 67:1187–1197.
- Mosconi MW, et al. (2009) Longitudinal study of amygdala volume and joint attention in 2- to 4-year-old children with autism. *Arch Gen Psychiatry* 66:509–516.
- Sparks BF, et al. (2002) Brain structural abnormalities in young children with autism spectrum disorder. *Neurology* 59:184–192.
- Nordahl CW, et al. (2012) Increased rate of amygdala growth in children aged 2 to 4 years with autism spectrum disorders: A longitudinal study. *Arch Gen Psychiatry* 69:53–61.
- Dalton KM, et al. (2005) Gaze fixation and the neural circuitry of face processing in autism. *Nat Neurosci* 8:519–526.
- Kliemann D, Dziobek I, Hatri A, Baudewig J, Heekeren HR (2012) The role of the amygdala in atypical gaze on emotional faces in autism spectrum disorders. *J Neurosci* 32:9469–9476.
- Bauman M, Kemper TL (1985) Histoanatomic observations of the brain in early infantile autism. *Neurology* 35:866–874.
- Schumann CM, Amaral DG (2006) Stereological analysis of amygdala neuron number in autism. *J Neurosci* 26:7674–7679.
- Wegiel J, et al. (2014) Stereological study of the neuronal number and volume of 38 brain subdivisions of subjects diagnosed with autism reveals significant alterations restricted to the striatum, amygdala and cerebellum. *Acta Neuropathol Commun* 2:141.
- Zhang XM, et al. (2009) Doublecortin-expressing cells persist in the associative cerebral cortex and amygdala in aged nonhuman primates. *Front Neuroanat* 3:17.
- Chareyron LJ, Lavenex PB, Lavenex P (2012) Postnatal development of the amygdala: A stereological study in rats. *J Comp Neurol* 520:3745–3763.
- Rubinow MJ, Juraska JM (2009) Neuron and glia numbers in the basolateral nucleus of the amygdala from preweaning through old age in male and female rats: A stereological study. *J Comp Neurol* 512:717–725.
- Courchesne E, et al. (2011) Neuron number and size in prefrontal cortex of children with autism. *JAMA* 306:2001–2010.
- Amaral DG, Schumann CM, Nordahl CW (2008) Neuroanatomy of autism. *Trends Neurosci* 31:137–145.
- O'Rahilly R, Müller F (2008) Significant features in the early prenatal development of the human brain. *Ann Anat* 190:105–118.
- Lee AL, Ogle WO, Sapolsky RM (2002) Stress and depression: Possible links to neuron death in the hippocampus. *Bipolar Disord* 4:117–128.
- McEwen BS (2004) Protection and damage from acute and chronic stress: Allostasis and allostatic overload and relevance to the pathophysiology of psychiatric disorders. *Ann N Y Acad Sci* 1032:1–7.
- Green SA, et al. (2015) Neurobiology of sensory overresponsivity in youth with autism spectrum disorders. *JAMA Psychiatry* 72:778–786.
- Kleinhaus NM, Richards T, Greenon J, Dawson G, Aylward E (2016) Altered dynamics of the fMRI response to faces in individuals with autism. *J Autism Dev Disord* 46:232–241.
- Cunningham CL, Martínez-Cerdeño V, Noctor SC (2013) Microglia regulate the number of neural precursor cells in the developing cerebral cortex. *J Neurosci* 33:4216–4233.
- Morgan JT, et al. (2010) Microglial activation and increased microglial density observed in the dorsolateral prefrontal cortex in autism. *Biol Psychiatry* 68:368–376.
- Morgan JT, Barger N, Amaral DG, Schumann CM (2014) Stereological study of amygdala glial populations in adolescents and adults with autism spectrum disorder. *PLoS One* 9:e110356.
- van Kooten IAJ, et al. (2008) Neurons in the fusiform gyrus are fewer and smaller in autism. *Brain* 131:987–999.
- Kennedy DP, Adolphs R (2012) The social brain in psychiatric and neurological disorders. *Trends Cogn Sci* 16:559–572.
- Berretta S, Pantazopoulos H, Lange N (2007) Neuron numbers and volume of the amygdala in subjects diagnosed with bipolar disorder or schizophrenia. *Biol Psychiatry* 62:884–893.
- Barger N, Sheley MF, Schumann CM (2015) Stereological study of pyramidal neurons in the human superior temporal gyrus from childhood to adulthood. *J Comp Neurol* 523:1054–1072.
- Heinsen H, Arzberger T, Schmitz C (2000) Celloidin mounting (embedding without infiltration)—A new, simple and reliable method for producing serial sections of high thickness through complete human brains and its application to stereological and immunohistochemical investigations. *J Chem Neuroanat* 20:49–59.
- Schumann CM, Amaral DG (2005) Stereological estimation of the number of neurons in the human amygdaloid complex. *J Comp Neurol* 491:320–329.
- West MJ (1993) New stereological methods for counting neurons. *Neurobiol Aging* 14:275–285.
- Lord C, Rutter M, Le Couteur A (1994) Autism Diagnostic Interview-Revised: A revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders. *J Autism Dev Disord* 24:659–685.
- Gundersen HJG, et al. (1988) The new stereological tools: Disector, fractionator, nucleator and point sampled intercepts and their use in pathological research and diagnosis. *APMIS* 96:857–881.